

Cytogenotoxic effect of trichothecene T2 toxin on *Allium sativum* root tip meristematic cells

Nasreen Jalal Hussein¹ and Asia A. M. Saadullah^{2*}

¹Department of Biology/ College of Science/ University of Zakho, Zakho, Kurdistan region/Iraq; nasreen.hussein@uoz.edu.krd . <https://orcid.org/0000-0002-2221-1330>.

²Department of Biology/ College of Science/ University of Duhok, Duhok, Kurdistan region/ Iraq; asia.saadullah@uod.ac . ORCID: <https://orcid.org/0000-0001-6166-4394>

* Correspondence: asia.saadullah@uod.ac

Available from. <http://dx.doi.org/10.21931/RB/2024.09.01.45>

ABSTRACT

Trichothecene T2 is a mycotoxin from the *Fusarium* species. This research aims to test the effect of the Trichothecene T2 toxin on mitotic index% (M.I.%) and induction of mitotic aberrations by using the *Allium sativum* (garlic) test system. The toxin concentrations in ppm were: 0.00, 0.30, 0.60, 0.90, and 1.20 for 12 hours. The garlic roots were then cut and mitotic slides were prepared by the squash method and examined under a light microscope. The results revealed that the mycotoxin has a significant mitodepressive effect at all concentrations compared to the control and the MI% reduction was proportional to increasing toxin concentration. The highest reduction in mitotic index was observed in the 1.2 ppm treatment. Moreover, this mycotoxin induced and increased the rate of mitotic abnormalities% (MA%) with increasing the mycotoxin concentration. The observed mitotic abnormalities were star-shaped anaphase, sticky metaphase, C-mitosis, sticky anaphase, depolarization, micronuclei, laggard chromosomes, anaphase bridges, and chromosome loss. The least frequently observed abnormality was micronuclei compared to the most frequent aberration, laggard chromosomes. The total mitotic abnormalities significantly increased with increasing the toxin dose concentration. These results suggest that this mycotoxin can inhibit the mitotic activity of the meristematic cells, it is mutagenic and can disrupt the spindle fibers activity of the dividing cells at all concentrations, especially at its higher doses in food. Therefore, the foods must be tested for fungi producing this mycotoxin.

Keywords: Mycotoxin; mitodepressive; root tip; spindle fibers; mutagenic

INTRODUCTION

Mycotoxins are natural chemicals produced by fungus that infect cereals and other agricultural foods. They have a detrimental impact on Animal and human health. They are also thermally stable and easily transmitted to humans through animal products, sophisticated food preparation, and food chains¹: toxic compounds are produce in foods and feeds by several strains of toxigenic fungi², which ultimately cause health problems in humans and animals³. Trichothecenes (TCTs) are a family of chemically related mycotoxins produced most commonly by filamentous fungi such as *Fusarium*, *Myrothecium*, *Stachybotrys*, *Trichoderma*, and *Trichothecium*, *Verticimonosporium*, *Cylindrocarpon*, threatening the health of both animals and human^{4,5}. The fungi that produce TCT are found all over the world. Their ability to grow under a variety of environmental conditions, including nutrient content, moisture content, temperature, and oxygen level in growth medium, results in successful colonization^{6,7}. T-2 toxin can be found in diverse regions worldwide and affects negatively on

both human and animal health. It affects many organs and organ systems in the various ways: neurotoxicity, immunotoxicity, hepatotoxicity, dermal toxicity, nephrotoxicity, as well as the reproductive system disruption, maternotoxic and embryo lethal⁸. T-2 toxin is thought to be a primary contributor in human alimentary toxic aleukia. TCT are non-volatile sesquiterpenoids with low molecular weight (usually 200-500 Da) produced by the terpenoid biosynthesis pathway^{9,10}. They are somewhat soluble in water but very soluble in polar organic solvents such ethyl acetate, chloroform, ethanol, methanol, and propylene glycol¹. TCTs share a three-ring molecule called 12,13-epoxytrichothec-9-ene (EPT)^{11,12}. The A-ring cyclohexene is fused to the B-ring tetrahydropyran, which is bridged by a two-carbon chain at C-2 and C-5, generating a cyclopentyl moiety (C-ring)¹³. TCT are classified into four kinds (A-D) based on the substitution pattern of EPT¹⁴. A hydroxyl (OH) group distinguishes type A TCT.

Define C-8 as having a group, an ester function, or no oxygen substitution¹⁵. Trichothecenes are quickly absorbed and suppress protein synthesis in growing tissues. Mycotoxins are natural chemicals generated by fungus that infect grains and other agricultural goods. Trichothecenes mycotoxin produced by the species *Fusarium* and *Trichoderma* is becoming increasingly important in agriculture globally because to the potential health implications. The most dangerous type A trichothecene mycotoxin is T-2 toxin, an unavoidable environmental hazard. T-2 toxin has a considerable detrimental effect on reproduction. Plant systems like onion, garlic and bean are widely used to determine the cytotoxic and genotoxic effects of various chemicals like food additives, and various pesticides, and different toxins^{16,17,18}. Trichothecenes are important foodborne mycotoxins implicated in human health and have immune cytotoxic effects. Toxicity data on HT-2 toxin are very limited. To the best of the authors' knowledge, there is no published data that tests the cytogenotoxic effects of this mycotoxin on *A. sativum* root tip meristematic cells. Therefore, the aim of this study is to test the Cytogenotoxic effect of T-2 mycotoxin on mitotic index and T-2 toxin-induced mitotic aberrations in *A. sativum* dividing cells at the root apical meristem.

MATERIALS AND METHODS

Garlic heads (*Allium sativum*, 2n=16) were purchased from a local store in Zakho, free from fertilizers.

Source of the mycotoxin T2 toxin:

Mycotoxin used in the current study is the 8230 Veratox for T-2/HT-2 (NEOGEN company /USA), Cross Reactivity T-2 toxin 100%, HT-2 toxin B 100%. The product details is Veratox® for T-2/HT-2 is a competitive direct ELISA that provides a quantitative analysis of T-2 toxin and HT-2 toxin in commodities such as wheat, rye, barley, oats, and corn. The different concentrations were diluted by using distilled water.

The garlics were placed on tubes filled with distilled water for a few days until the roots were emerged and reached the length of about 1 cm then they were removed from those tubes and placed on another set of tubes with different concentrations of the mycotoxin T2 for about 12 hours. Then the roots were harvested and prepared microscopic slides for the root tip by squash method and stained with Feulgen Giemsa stain and visualized under light microscopy¹⁹.

The mitotic index calculation:

The MI% calculation was done according to the following formula as published by Fiskesjö, (1985).

$$\text{M.I.\%} = \text{DC/TC} * 100 \quad (1)$$

Where, M.I.= Mitotic Index, DC=Dividing Cells, TC=Total counted Cells.

Mitotic abnormalities percentage:

The mitotic abnormalities percentage was calculated by the following formula (Palsikowski et al., 2017)

$$\text{MA\%} = \text{ADC/TC} * 100 \quad (2)$$

Where, MA=Mitotic Abnormalities, ADC=Abnormal dividing cells, TC=Total counted cells.

Statistical analysis:

Most of the statistics were done using SPSS (Statistical Package for the Social Sciences) software version 14, by One way ANOVA, and differences between all the treatments were determined by Duncan's multiple-range test²⁰. The data were calculated as mean \pm standard error. Each experiment was repeated at least three times. The significance was measured at $P < 0.05$. The bar graphs were generated using GraphPad Prism version 9.0.

RESULTS

Trichothecenes are a class of structurally related mycotoxins that have different degrees of cytotoxicity. Because of the potential health risks, trichothecene (TCT) mycotoxin is becoming increasingly important in agriculture around the globe. It is primarily metabolized and eliminated after consumption, producing more than 20 metabolites, the most important of which is the hydroxy trichothecenes-2 toxin. (A12). The effect of T2 toxin on mitotic index% is shown in Table 1 and figure, 1.

Treatments (ppm)	Time (hours)	Total no. of cells	interphase	Dividing cells	Mitotic index%
0.00	12	1127.000 \pm 23.643	1009.400 \pm 19.954 a, b	117.600 \pm 4.345 a	10.4300 \pm 0.222 a
0.30	12	990.000 \pm 45.655	915.928 \pm 42.771 b	74.072 \pm 2.942 b	7.4867 \pm 0.075 b
0.60	12	1102.000 \pm 56.580	1049.593 \pm 53.323 a	52.407 \pm 3.311 c	4.75 \pm 0.078 c
0.90	12	1052.333 \pm 26.194	1013.816 \pm 25.189 a, b	38.518 \pm 1.073 d	3.68 \pm 0.036 d
1.20	12	996.667 \pm 3.844	978.223 \pm 4.019 a, b	18.444 \pm 0.939 e	1.851 \pm 0.095 e
Significance			n.s.	***	***

¹ Note: n.s. means non-significant, *** means significant at $p < 0.001$. Different superscript letters were produced by DMRT.

Table 1: The effect of T2 toxin on the mitotic index% of *A. sativum* root tip cells.




Figure 1: The effect of T2 toxin on no. of cells in interphase and dividing cells, and mitotic index% of *A. sativum* root tip cells.

The highest concentration, 1.5 ppm which caused it to decrease from 10.53 in the control to 0.7633 at 1.5 ppm. The mitodepressive effect was significant at $P < 0.001$.




Figure 2: Normal mitosis and different abnormalities in *A sativum* root apical meristematic cells: a-normal interphase, b- normal prophase, c- normal metaphase, d- normal anaphase, e- normal telophase, f- Star-shaped anaphase, g- Sticky Metaphase, h- C-Mitosis, i- Sticky anaphase, j- Depolarization, k- Laggard Chromosomes, l- Anaphase Bridge, m- Micronucleus, n- Chromosome Loss.

The effect of the T2 toxin on chromosomal aberrations% is shown in table, 2 and figure, 3. According to the table, the aberrations increased proportionally with increasing mycotoxin concentration. Total mitotic abnormalities increased proportionally with increasing the concentration of the mycotoxin. The observed mitotic abnormalities were star-shaped anaphase, sticky metaphase, C-mitosis, sticky anaphase, depolarization, micronuclei, laggard chromosomes anaphase bridges, and chromosome loss. The least frequently observed abnormality was micronuclei at 0.00 in the control and increased to 0.1672 in the 1.2 ppm of T2. As compared to the most frequent aberration, laggard chromosomes were 0.0536% in the control and increased to 1.1341% in 1.2 ppm of the tested mycotoxin. The total CAI% were 0.1106 in the control group, which increased to about 5.2168% at the highest concentration of the mycotoxin. There were significant differences between treatments at $p < 0.001$ for all mitotic aberrations.

T2 Concentration (ppm)	Star shaped anaphase%	Sticky metaphase%	C-Mitosis%	Sticky anaphase%	Depolarization%	Laggard Chromosomes%	Anaphase bridges%	Micronuclei%	Chromosome loss%	Total CA %
0.00	0.0536	0.0536	0.0536	0.0536	0.0536	0.0536	0.0536	0.00	0.0536	0.1106
1.2	1.1341	1.1341	1.1341	1.1341	1.1341	1.1341	1.1341	0.1672	1.1341	5.2168

0.0	0.000 ± 0.000 d	0.0000 ± 0.000 d	0.0571 ± 0.022 c	0.0000 ± 0.000 e	0.000 ± 0.000 d	0.0536 ± 0.0192 5 e	0.000 ± 0.000 c	0.000 ± 0.000 c	0.000 ± 0.000 e	0.1106 ± 0.03676 d
0.3	0.0761 ± 0.011 c	0.1175 ± 0.008 c	0.5833 ± 0.039 b	0.1910 ± 0.027 d	0.245 ± 0.030 c	0.2406 ±0.035 18 d	0.004 ± 0.004 c	0.000 ± 0.000 c	0.108 ± 0.003 d	1.5644 ±0.0815 8 c
0.6	0.100 ± 0.005 c	0.4700 ± 0.041 a	1.0310 ± 0.058 a	0.3741 ± 0.012 c	0.287 ± 0.025 b, c	0.4142 ± 0.0029 0 c	0.079 ± 0.008 b	0.0000 ± 0.000 c	0.174 ± 0.0120 c	2.9284 ±0.1397 5 b
0.9	0.1869 ± 0.008 b	0.2466 ± 0.027 b	0.6432 ± 0.005 b	0.5285 ± 0.019 b	0.358 ± 0.016 b	0.6268 ± 0.0201 9 b	0.085 ± 0.0074 2 b	0.1112 ± 0.013 b	0.387 ± 0.004 b	3.1728 ±0.0207 7 b
1.2	0.2802 ± 0.010 a	0.4619 ± 0.031 a	0.9601 ± 0.033 a	0.8026 ± 0.015 a	0.532 ± 0.034 a	1.1341 ± 0.0397 2 a	0.197 ± 0.010 a	0.1672 ± 0.016 a	0.682 ± 0.039 a	5.2168 ±0.0526 3 a
F value	192.205	62.803	116.06 4	329.04 3	63.116	240.15 3	142.99 0	70.784	212.055	591.802
Significance	***	***	***	***	***	***	***	***	***	***

*** means significant at $p < 0.001$. superscript letters are produced by DMRT

Table 2: the effect of T2 toxin on mitotic abnormalities% of *A. sativum* root tip meristematic cells after 12 hours.

The abnormalities caused by this mycotoxin in *A. sativum* root tip meristematic cells were: star-shaped anaphase, sticky metaphase, C-Mitosis, sticky anaphase, depolarization, micronuclei, and laggard Chromosomes. The most frequent abnormality was C-mitosis and sticky anaphase. According to Table, 2, the total abnormalities were increasing with increasing the toxin concentration. The highest number was observed with the 1.5 ppm treatment. The different abnormalities along with normal mitosis are shown in Figure, 2. The mycotoxin T2 induces the distortion and inhibition of mitotic spindle formation which causes the C-mitotic cells and stickiness in metaphase and anaphase. Moreover, the induction of chromosome loss and micronuclei.




Figure 3: The effect of T2 toxin on mitotic abnormalities in *A. sativum* root tip meristematic cells.

About 1000 cells were counted for each slide and each experiment was repeated at least three times. According to table 1, the different concentrations of the mycotoxin caused a significant reduction in the mitotic index at all treatments along with reduction in the dividing cells number. However, the greatest effect on the mitotic index was observed at the highest concentration of the mycotoxin treatment (1.2 ppm) which caused the

mitotic index% to decrease from (10.4300 ± 0.222) in the control group to (1.851 ± 0.095) at 1.2 ppm treatment. The mitodepressive effect was significant at $P < 0.001$. Moreover, there were significant differences among the treatments as detected by DMRT.

The effect of the T2 toxin on chromosomal aberrations% is shown in table, 2 and figure, 3. The abnormalities caused by this mycotoxin in *A. sativum* root tip meristematic cells were: star-shaped anaphase, sticky metaphase, C-Mitosis, sticky anaphase, depolarization, micronuclei, and laggard chromosomes. The most frequent abnormalities were C-mitosis and sticky anaphase. According to Table, 2, the total abnormalities were proportionally increasing with increasing toxin concentration. The highest number was observed with the 1.2 ppm treatment. The different abnormalities along with normal mitosis are shown in Figure, 2.

The total MA% was (0.1106 ± 0.03676) in the control group, which increased to about (5.2168 ± 0.05263) at the highest concentration of the mycotoxin used. There were significant differences between treatments at $p < 0.001$ for all mitotic aberrations. The ability to induce different abnormalities in the *A. sativum* cells at the root apical meristem indicates its mutagenic activity.

DISCUSSION

According to table, 1 the different concentrations of mycotoxin caused a significant reduction in the mitotic index at all treatments. However, the greatest reduction in the mitotic index was observed at. other research which tested the effect of Aflatoxin B1 induces reactive oxygen species-dependent caspase-mediated apoptosis in normal human cells, inhibits *Allium cepa* root cell division²¹, and another research tested the toxic activity of citrinin, a fungal phytotoxin, and its mode of action in onion cells²².

The abnormalities caused by this mycotoxin in *A. sativum* root tip meristematic cells were: star-shaped anaphase, sticky metaphase, C-Mitosis, sticky anaphase, depolarization, micronuclei, and laggard Chromosomes. The most frequent abnormality was C-mitosis and sticky anaphase. According to Table, 2, the total abnormalities were increasing with increasing the toxin concentration. The highest number was observed with the 1.5 ppm treatment. The different abnormalities along with normal mitosis are shown in Figure, 2. The mycotoxin T2 induces the distortion and inhibition of mitotic spindle formation which causes the C-mitotic cells and stickiness in metaphase and anaphase. Moreover, the induction of chromosome loss and micronuclei.

The ability to induce different abnormalities in the *A. sativum* cells at the root apical meristem indicates its mutagenic activity. The induction of both Micronuclei and stickiness are the most obvious indicators of cytotoxicity. According to Table 2, the increase in the frequency of micronuclei was dose-dependent. These results are in agreement with results obtained by²³. The total abnormality percentage increase was also dose-dependent. This increase is in agreement with previous results of testing various substances on onion root tips^{24,25}.

These results are in agreement with other researches which tested the effect of Aflatoxin B1 on reactive oxygen species-dependent caspase-mediated apoptosis in normal human cells, and on inhibition of *Allium cepa* root cell division²⁶, and another research tested the toxic activity of citrinin, a fungal phytotoxin, and its mode of action in onion cells²⁷. The mycotoxin T2 induces the distortion and inhibition of mitotic spindle formation which causes the C-mitotic cells and stickiness in metaphase and anaphase. Moreover, the induction of chromosome loss and micronuclei. The least frequently observed abnormality was micronuclei at (0.000 ± 0.000) in the control and increased to (0.1672 ± 0.016) in the 1.2 ppm of T2. On the other hand, the most frequent aberration observed was laggard chromosomes were (0.0536 ± 0.01925) in the control and increased to (1.1341 ± 0.03972) in 1.2 ppm of the tested mycotoxin. A laggard chromosome is defined as the chromosome that did not overlap along the long axis of the spindle with any of the properly segregating chromosomes²⁸. The induction of both Micronuclei and stickiness are the most obvious indicators of cytotoxicity, Micronuclei are small membrane bounded compartments with a DNA content encapsulated by a nuclear envelope and spatially separated from the primary nucleus (Krupina et al., 2021). Chromosome stickiness may result from chromatin fibers' sticking to each other or breaking due to erroneous or inadequate condensation of these fibers, as a consequence of this, movement of mitotic spindle fibers together with inner-chromosome stickiness when the chromosome is drawn to the pole causes secondary anomalies (bridge and fragment occurrence), Stickiness in chromosomes is an indication of the high toxicity of the chemical substance and usually this may kill the cells with the irreversible damages²⁵

CONCLUSIONS

The mycotoxin T2 is a mutagen and it inhibits cell division and induce a wide range of abnormalities in *A sativum* root apical meristematic cells at lower doses and is very strong mutagen. Therefore, the food grains that contain this mycotoxin should be tested carefully to ensure that they are not contaminated with the T2 toxin producing fungi to avoid its hazardous health risks.

REFERENCES

1. Polak-Śliwińska, M.; Paszczyk, B. 2021. Trichothecenes in Food and Feed, Relevance to Human and Animal Health and Methods of Detection: A Systematic Review. *Molecules*. 26, 454.
2. Bryden, W. L. 2007. Mycotoxins in the food chain: Human health implications. *Asia Pacific journal of clinical nutrition*. 16:95–101.
3. Bennett, J. W., and Klich, M. 2003. Mycotoxins. *Clinical Microbiology Reviews*. 16:497–516.
4. Chen, P.; Xiang, B.; Shi, H.; Yu, P.; Song, Y.; Li, S. Recent advances on type A trichothecenes in food and feed: Analysis, prevalence, toxicity, and decontamination techniques. *Food Control*. 2020, 118, 107371.
5. Milićević, D.R.; Škrinjar, M.; Baltić, T. 2010. Real and Perceived Risks for Mycotoxin Contamination in Foods and Feeds: Challenges for Food Safety Control. *Toxins*. 2, 572.
6. McCormick, S.P.; Stanley, A.M.; Stover, N.A.; Alexander, N.J. 2011. Trichothecenes: From simple to complex mycotoxins. *Toxins*. 3, 802–814.
7. Foroud, N.A.; Baines, D.; Gagkaeva, T.Y.; Thakor, N.; Badea, A.; Steiner, B.; Bürstmayr, M.; Bürstmayr, H. 2019. Trichothecenes in Cereal Grains—An Update. *Toxins*. 11, 634.
8. S K Kudury, I A Abed and B A Mahdi . The Effect of Local Bio Fertilizer and Their Enzymatic Activity on Growth of Maize Plant and some Biological Therites in the Soil. *IOP Conf Ser Earth Environ Sci* 2023, 1252 (1), 012050. <https://doi.org/10.1088/1755-1315/1252/1/012050>.
9. Arunachalam, C., and Doohan, F.M. 2013. Trichothecene toxicity in eukaryotes: Cellular and molecular mechanisms in plants and animals. *Toxicology Letters*. 217, 149–158.
10. Bottalico, A., & Perrone, G. (2002). Toxicogenic *Fusarium* species and mycotoxins associated with head blight in small-grain cereals in Europe. *Mycotoxins in Plant Disease: Under the aegis of COST Action 835 ‘Agriculturally Important Toxicogenic Fungi 1998-2003’*, EU project (QLK 1-CT-1998-01380), and ISPP ‘*Fusarium* Committee’, 611-624.
11. Çavuşoğlu, D. 2022. Powerful toxic activity of citrinin, a fungal phytotoxin, and its mode of action in onion cells. *Environmental Science and Pollution Research*, 29(4): 6205-6218.
12. A A Al-Azzami , Th T Mohammed . Effect of Adding Dry Leaves of Lemongrass (*Cymbopogon Citratus*) To the Diet on Some Biochemical Tests of Blood in Broiler (Ross 308). *IOP Conf Ser Earth Environ Sci* 2023, 1252 (1), 12125. <https://doi.org/10.1088/1755-1315/1252/1/012125>.
13. Das, T., S. Hazra, S. Sengupta, P. Hazra, and D. Chattopadhyay. 2021. Genotoxic effect of saccharin on *Allium cepa* root tips. *Biologia*. 76: 3191-3199.
14. R S Obaid, S H S Al-Warshan and I A Abed . Efficiency of Some Clays and Organic Materials on the Reduction of Aflatoxin B1 Produced from Isolates of the Fungus *Aspergillus flavus* Contaminating Corn Grains. *IOP Conf Ser Earth Environ Sci* 2023, 1252 (1), 012003. <https://doi.org/10.1088/1755-1315/1252/1/012003>.
15. Duncan, D. B. 1955. Multiple range and multiple F tests. *Biometrics*. 11: 1-42.
16. EFSA (European Food Safety Authority) Scientific Opinion on the risks for animal and public health related to the presence of T-2 and HT-2 toxin in food and feed. *EFSA Journal*. 2011:2481–2668. doi: 10.2903/j.efsa.2011.2481.
17. A A Al-Azzami , Th T Mohammed . The Effect of Adding Lemongrass Leaf Powder (*Cymbopogon Citratus*) to the Diet as a Natural Supplement on Some Productive Traits and Oxidation Indicators in Broiler (Ross 308). *IOP Conf Ser Earth Environ Sci* 2023, 1252 (1), 12123. <https://doi.org/10.1088/1755-1315/1252/1/012123>.

18. Foroud, N.A.; Shank, R.A.; Kiss, D.; Eudes, F.; Hazendonk, P. 2016. Solvent and Water Mediated Structural Variations in Deoxynivalenol and Their Potential Implications on the Disruption of Ribosomal Function. *Frontiers in Microbiology*. 7, 1239.
19. Garvey, G.S.; McCormick, S.P.; Rayment, I. Structural and Functional Characterization of the TRI101 Trichothecene 3-O-Acetyltransferase from *Fusarium sporotrichioides* and *Fusarium graminearum*: Kinetic insights to combating fusarium head blight. *Journal of Biological Chemistry*, 283(3), 1660-1669.
20. Hussein, N. J. 2023. "Evaluation of Cytogenotoxic Effect of Potassium Acetate on *Allium Cepa* L. Root Tips". *Emirates Journal of Food and Agriculture*. 35(8):708-714. doi:10.9755/ejfa.2023.3129.
21. Ameen M. Shaman , Th. T. Mohammed. Effect Using Feed Additives Instead of Imported Premixes Affects the Physiology of Broiler Chickens. *IOP Conf Ser Earth Environ Sci* 2023, 1262 (7), 72080. <https://doi.org/10.1088/1755-1315/1262/7/072080>.
22. Jebur, S. F.; Abdulateef, S. M. Effect of The Critical Period and Gene Silencing on Blood Cellular Traits in Local Chicken and Level of Welfare. *IOP Conf Ser Earth Environ Sci* 2023, 1252 (1), 12141. <https://doi.org/10.1088/1755-1315/1252/1/012141> .
23. Nielsen, C.; Casteel, M.; Didier, A.; Dietrich, R.; Märtlbauer, E. 2009. Trichothecene-induced cytotoxicity on human cell lines. *Mycotoxin Research*. 25, 77–84.
24. Palsikowski, P. A., Roberto, M. M., Sommaggio, L. R., Souza, P. M., Morales, A. R., & Marin-Morales, M. A. (2018). Ecotoxicity evaluation of the biodegradable polymers PLA, PBAT and its blends using *Allium cepa* as test organism. *Journal of Polymers and the Environment*. 26: 938-945.
25. Pestka, J. J. 2007. Deoxynivalenol: Toxicity, mechanisms and animal health risks. *Animal Feed Science and technology*. 137: 283–298.
26. Elkhateeb, S. Z.; Ebraheem, M. O.; Abdulateef, S. M.; Ahmed, I. A. Constraints Affecting the Welfare of Domestic Sheep Grazing in the Natural Pasture. *IOP Conf Ser Earth Environ Sci* 2023, 1252 (1), 12144. <https://doi.org/10.1088/1755-1315/1252/1/012144>.
27. Saadullah, A. A. M. (2022). Studies on teratogenic and maternal effects of Trichothecene (TCT) extracted from *Fusarium* and *Trichoderma* culture on pregnant Albino Mice. *Revis Bionatura*. 2023; 8 (2) 58.
28. Sharma, A. and A. Sharma. 1999. *Chromosome Techniques*. 3rd ed. Buterworths, London.

Received: October 9th 2023/ **Accepted:** January 15th 2024 / **Published:** 15 February 2024

Citation: Hussein N. J. and Saadullah A. A. M. Cytogenotoxic effect of trichothecene T2 toxin on *Allium sativum* root tip meristematic cells. *Revis Bionatura* 2024; 9 (1) 45.
<http://dx.doi.org/10.21931/RB/2024.09.01.45>

Additional information Correspondence should be addressed to asia.saadullah@uod.ac

Peer review information. Bionatura thanks anonymous reviewer(s) for their contribution to the peer review of this work using <https://reviewerlocator.webofscience.com/>

All articles published by Bionatura Journal are made freely and permanently accessible online immediately upon publication, without subscription charges or registration barriers.

Bionatura ISSN. First 13909355 Ecuador. **Scopus coverage years:** from 2016 to the present

Publisher's Note: Bionatura stays neutral concerning jurisdictional claims in published maps and institutional affiliations.

Copyright: © 2023 by the authors. They were submitted for possible open-access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

